

Soluciones a algunos problemas gráficos correspondientes a los problemas de Matemáticas II para 2º de Bachillerato


**Pedro González Ruiz
Catedrático de Matemáticas**

Versión 21: julio de 2024


Capítulo 2

Sección 2.2


6.




7.



8.




9.




Sección 2.8

7.





10.




Capítulo 4

Sección 4.1






[10]




[12]



[11]




[13]




[14]




[15]




[18]





[20]




[19]




[21]




[22]




[23]




[24]



[25]




[26]




Sección 4.2


1.




2.




4.




5.




6.




7.




8.




9.




10.




11.




14.



20.




21.




Capítulo 6

3.




$$\begin{aligned}
 S &= \int_1^3 |x^2 - 4x + 3| dx = \\
 &= \int_1^3 -(x^2 - 4x + 3) dx = \frac{4}{3}
 \end{aligned}$$

4.



$$S = \int_0^2 [(-x^2 + 4x) - 2x] dx = \frac{4}{3}$$

5.


Respecto al eje X :

$$\begin{aligned} S &= \int_1^2 (x^2 - 1) dx + \int_2^8 \left(\frac{8}{x} - 1 \right) dx = \\ &= \frac{4}{3} + (16 \ln 2 - 6) = 16 \ln 2 - \frac{14}{3} \end{aligned}$$

Más fácil con respecto al eje Y :


$$S = \int_1^4 \left(\frac{8}{y} - \sqrt{y} \right) dy = 16 \ln 2 - \frac{14}{3}$$

7.




$$S = \int_2^{2\sqrt{3}} \frac{1}{4+x^2} dx = \frac{\pi}{24}$$

8.




$$\begin{aligned}
 S &= \int_{-\frac{9}{2}}^{-4} -\ln(x+5) dx + \int_{-4}^1 \ln(x+5) dx = \\
 &= \left(-\frac{1}{2} \ln 2 + \frac{1}{2} \right) + (6 \ln 6 - 5) = \\
 &= 6 \ln 6 - \frac{1}{2} \ln 2 - \frac{9}{2}
 \end{aligned}$$

9.




$$S = \int_1^e \ln x dx = 1$$

10.




$$S = \int_0^4 [(6x - x^2) - (x^2 - 2x)] dx = \frac{64}{3}$$

12.




$$S = \int_0^3 [(2x - x^2) + x] dx = \frac{9}{2}$$

14.




$$S = \int_1^5 [(2x - 5) - (x^2 - 4x)] dx = \frac{32}{3}$$

18.




$$\begin{aligned} S &= \int_0^2 (x^3 - 6x^2 + 8x) dx + \int_2^4 -(x^3 - 6x^2 + 8x) dx = \\ &= 4 + 4 = 8 \end{aligned}$$


20.




21.



22.



23.




Sea S_1 la superficie de la región rayada:

$$S_1 = \int_0^1 [(3-x) - (1+x^5)] dx = \frac{4}{3}$$

Sea S_2 la superficie de la región sombreada:


$$S_2 = \frac{9}{2} - \frac{4}{3} = \frac{19}{6}$$

24.



$$S = \int_0^1 (\sqrt{x} - x^2) dx = \frac{1}{3}$$

26.




La recta tangente en $x = e$ es:

$$y = 3ex - 2e^2$$

La superficie del triángulo de vértices $(2e/3, 0)$, $(e, 0)$, (e, e^2) es $e^3/6$, luego

$$S = \int_1^e x^2 \ln x dx - \frac{e^3}{6} = \frac{e^3 + 2}{18}$$


30.



La recta tangente en $x = 0$ es $y = 4x$.
La región es simétrica respecto de la recta $x = 2$ (demuéstrelo), luego:

$$S = 2 \int_0^2 [4x - (4x - x^2)] dx = \frac{16}{3}$$


31.



La recta tangente en $x = 1$ es $y = x - 1$.


$$S = \int_1^3 [(x - 1) - \ln x] dx = 4 - 3 \ln 3$$

32.




$$\begin{aligned} S &= \int_1^2 [\sqrt{x} + (x - 2)] dx + \\ &+ \int_2^4 [\sqrt{x} - (x - 2)] dx = \frac{13}{6} \end{aligned}$$

39.




$$\begin{aligned}
 S &= \int_0^{\pi/4} (\cos x - \sin x) dx + \\
 &\quad + \int_{\pi/4}^{\pi/3} (\sin x - \cos x) dx = \\
 &= (\sqrt{2} - 1) + \left(\sqrt{2} - \frac{1}{2} - \frac{\sqrt{3}}{2} \right) = \\
 &= 2\sqrt{2} - \frac{3}{2} - \frac{\sqrt{3}}{2}
 \end{aligned}$$

40.




$$S = \int_0^1 \left[(3x + 2) - \frac{2}{1+x^2} \right] dx = \frac{7-\pi}{2}$$

44.




$$\begin{aligned}
 S &= \int_{-1}^0 [(x^3 - 2x) - x^2] dx + \\
 &\quad + \int_0^2 [x^2 - (x^3 - 2x)] dx = \\
 &= \frac{5}{12} + \frac{8}{3} = \frac{37}{12}
 \end{aligned}$$

46.




$$S = \int_{-4}^{-1} [-x - (x+2)^2] dx = \frac{9}{2}$$

47.




$$\begin{aligned} S &= \int_{-4}^1 [(-x^2 - 3x + 10) - (x^2 + 3x + 2)] dx = \\ &= \frac{125}{3} \end{aligned}$$

51.



$$\begin{aligned} S &= \int_0^1 (x^2 + 1) dx + \int_1^2 \left[\frac{2}{x} - (x - 1) \right] dx = \\ &= \frac{4}{3} + \left(-\frac{1}{2} + 2 \ln 2 \right) = \frac{5}{6} + 2 \ln 2 \end{aligned}$$


55.



Por simetría:


$$S = 2 \int_0^\pi (2 \sin x - \sin 2x) dx = 8$$

57.




$$\begin{aligned} S &= \int_{-2}^{-1} (5x + 10) dx + \int_{-1}^3 (x^2 - 2x + 2) dx = \\ &= \frac{5}{2} + \frac{28}{3} = \frac{71}{6} \end{aligned}$$

62.




$$\begin{aligned} S &= \int_0^1 4x dx + \int_1^3 \frac{16}{(x+1)^2} dx + \int_3^4 (4-x) dx = \\ &= 2 + 4 + \frac{1}{2} = \frac{13}{2} \end{aligned}$$

63.




$$\begin{aligned} S &= \int_0^{2\pi/3} \left(\frac{1}{2} + \cos x \right) dx + \\ &\quad + \int_{2\pi/3}^{\pi} -\left(\frac{1}{2} + \cos x \right) dx = \\ &= \left(\frac{\pi}{3} + \frac{\sqrt{3}}{2} \right) + \left(-\frac{\pi}{6} + \frac{\sqrt{3}}{2} \right) = \\ &= \sqrt{3} + \frac{\pi}{6} \end{aligned}$$

64.




$$S = \int_{-1}^0 (e^{x+2} - e^{-x}) dx = (e - 1)^2$$

68.




$$S = \int_{-1}^0 -xe^{-x} dx = 1$$

71.



$$S = \int_0^1 [(1+x) - (1+x^2)] dx = \frac{1}{6}$$

72.




La recta tangente en $x = 3$ es:

$$y = \frac{ex}{3}$$


$$S = \int_0^3 \left(e^{x/3} - \frac{ex}{3} \right) dx = \frac{3e}{2} - 3$$

74.



$$\begin{aligned} S &= \int_0^2 \left(x^2 - \frac{x^2}{2} \right) dx + \int_2^4 \left(2x - \frac{x^2}{2} \right) dx = \\ &= \frac{4}{3} + \frac{8}{3} = 4 \end{aligned}$$

77.




La recta tangente en $x = 0$ es:

$$y = 1 - \frac{x}{2}$$

$$S = \int_0^2 \left[e^{-x} - \left(1 - \frac{x}{2} \right) \right] dx = 1 - \frac{2}{e}$$


82.



$$S_{\text{sombreada}} = \int_{-3}^{-1} [(x^3 + 3x^2) - (x + 3)] dx = 4$$


$$S_{\text{rayada}} = \int_{-1}^1 [(x + 3) - (x^3 + 3x^2)] dx = 4$$

83.



$$S = \int_0^2 x|x-2| dx = \int_0^2 -x(x-2) dx = \frac{4}{3}$$

84.




La recta tangente en $x = 0$ es:

$$y = x$$


$$S = \int_0^1 [x - \ln(1+x)] dx = \frac{3}{2} - 2\ln 2$$

86.




$$S = \int_{-\frac{1}{2}}^0 (e^{-2x} + 2ex) dx = \frac{e}{4} - \frac{1}{2}$$

88.




$$S = \int_{-1}^3 [(2x+2) - (x^2 - 1)] dx = \frac{32}{3}$$

89.




$$\begin{aligned}
 S &= \int_0^2 (x - x|x-1|) dx = \\
 &= \int_0^1 [x + x(x-1)] dx + \int_1^2 [x - x(x-1)] dx = \\
 &= \frac{1}{3} + \frac{2}{3} = 1
 \end{aligned}$$

90.



$$S = \int_{-1}^2 [2 - (x^3 - 3x)] dx = \frac{27}{4}$$

91.




$$S_{\text{sombreada}} = \int_0^{\sqrt{2}} \left(1 - \frac{x^2}{2}\right) dx = \frac{2\sqrt{2}}{3}$$

La superficie del rectángulo $ABCD$ es 2, luego


$$S_{\text{rayada}} = 2 - \frac{2\sqrt{2}}{3} = \frac{6 - 2\sqrt{2}}{3}$$

93.




$$S = \int_1^4 \left(5 - x - \frac{4}{x} \right) dx = \frac{15}{2} - 8 \ln 2$$

94.



$$S = \int_0^1 [x^2 + 4 - (2x + 3)] dx = \frac{1}{3}$$

95.




$$S = \int_0^{e-1} [1 - \ln(x + 1)] dx = e - 2$$

Muchísimo más fácil con respecto al eje Y:


$$S = \int_0^1 (e^y - 1) dy = e - 2$$

96.




$$S = \int_0^4 [(6x - x^2) - (x^3 - 2x)] dx = \frac{64}{3}$$

97.



$$\begin{aligned} S &= S_1 + S_2 = \int_{-2}^2 \left[(x + 5) - \left(4 - \frac{x^2}{4} \right) \right] dx + \\ &\quad \int_2^3 \left[(x + 5) - (x^2 - 1) \right] dx = \frac{16}{3} + \frac{13}{6} = \frac{15}{2} \end{aligned}$$

101.




Integrando respecto al eje X :

$$S = S_1 + S_2 = \int_1^3 \left(\frac{5-x}{2} - \frac{9-x^2}{4} \right) dx + \int_3^5 \frac{5-x}{2} dx = \frac{2}{3} + 1 = \frac{5}{3}$$

Integrando respecto al eje Y :


$$S = \int_0^2 \left[(5-2y) - \sqrt{9-4y} \right] dy = \frac{5}{3}$$

102.




$$\begin{aligned} S &= \int_{-1}^4 [x+4 - |x \cdot (x-2)|] dx = \\ &= \int_{-1}^0 [x+4 - x(x-2)] dx + \\ &\quad + \int_0^2 [x+4 + x(x-2)] dx + \\ &\quad + \int_2^4 [x+4 - x(x-2)] dx = \\ &= \frac{13}{6} + \frac{26}{3} + \frac{22}{3} = \frac{109}{6} \end{aligned}$$

103.




$$\begin{aligned} S &= \int_{3/2}^4 \left(-x^2 + 6x - 5 - \frac{x+2}{2} \right) dx = \\ &= \frac{125}{48} \end{aligned}$$

104.




$$S = 2 \int_0^1 \left(\frac{1}{1+x^2} - \frac{x}{2} \right) dx = \frac{\pi - 1}{2}$$

109.




Sea $f(x) = |x^2 - 4|$. Entonces: $S = 2(I_1 + I_2)$, $I_1 = \int_0^2 (5 - f(x)) dx = \frac{14}{3}$, $I_2 = \int_2^3 (5 - f(x)) dx = \frac{8}{3}$, y de aquí $S = \frac{44}{3}$.

110.




$$S = \int_1^3 (x - 1 - \ln x) dx = 4 - 3 \ln 3$$

112.




$$S = \int_0^2 [(-x^2 + 4x) - x^2] \, dx = \frac{8}{3}$$

114.




$$S = \int_0^1 x \, dx + \int_1^3 \frac{1}{x^3} \, dx = \frac{17}{18}$$

115.




$$\begin{aligned}
 S &= \int_0^4 [(6x - x^2) - |x^2 - 2x|] \, dx = \\
 &= \int_0^2 [(6x - x^2) - (2x - x^2)] \, dx + \\
 &\quad \int_2^4 [(6x - x^2) - (x^2 - 2x)] \, dx = \\
 &= 8 + \frac{32}{3} = \frac{56}{3}
 \end{aligned}$$

116.




$$\begin{aligned}
 S &= \int_1^2 [(4 - 2x) - (3 - x^2)] \, dx + \\
 &\quad \int_2^4 \left[(4 - 2x) - \left(-\frac{x^2}{4} \right) \right] \, dx = \\
 &= \frac{1}{3} + \frac{2}{3} = 1
 \end{aligned}$$

118.




$$\begin{aligned}
 S &= \int_{-\frac{1}{2}}^0 (e^{-2x} + 2ex) dx = \\
 &= \left[-\frac{1}{2}e^{-2x} + ex^2 \right]_{-\frac{1}{2}}^0 = \frac{e-2}{4}
 \end{aligned}$$

119.




$$\begin{aligned}
 S &= \int_1^3 \left[\ln(x+2) - \frac{x-3}{2} \right] dx = \\
 &= 5 \ln 5 - 3 \ln 3 - 1
 \end{aligned}$$

123.




$$S = 2 \int_0^2 [x - (x^2 - 2)] dx = \frac{20}{3}$$

124.




$$S = \int_0^4 (4x^3 - x^4) dx = \frac{256}{5}$$

127.




$$S = \int_{-2}^0 f(x) dx + \int_0^2 f(x) dx = 4 + \frac{8}{3} = \frac{20}{3}$$

129.




$$S = \int_0^1 (g(x) - f(x)) dx = \int_0^1 (-x^2 + 2x + 2 - x^3 - 2) dx = \left[-\frac{x^3}{3} + x^2 - \frac{x^4}{4} \right]_0^1 = \frac{5}{12}$$

133.



$$S = 2 \int_1^2 \left[(5 - x^2) - \frac{4}{x^2} \right] dx = 2 \left[5x - \frac{x^3}{3} + \frac{4}{x} \right]_1^2 = \frac{4}{3}$$

136.



$$\begin{aligned}
 S &= 2 \int_0^2 (f(x) - g(x)) dx = 2 \int_0^2 (-x^2 + 7 - |x^2 - 1|) dx = \\
 &= 2 \left[\int_0^1 (-x^2 + 7 - (1 - x^2)) dx + \int_1^2 (-x^2 + 7 - (x^2 - 1)) dx \right] = \frac{56}{3}
 \end{aligned}$$